

Baker-Beynon duality beyond semisimplicity

Luca Carai, Universitat de Barcelona

Joint work with: Serafina Lapenta and Luca Spada

Algebra Week
Università di Siena, 5 July 2023

Structure of the talk

Part I: Generalize Baker-Beynon duality to non-semisimple abelian ℓ -groups and Riesz spaces.

Part II: Non-standard analysis techniques to get a geometrical understanding of the dual objects.

Part I

A generalization of Baker-Beynon duality

Baker-Beynon duality

Definition

- An **(abelian) ℓ -group** is an abelian group A equipped with a lattice order such that $a \leq b$ implies $a + c \leq b + c$ for every $a, b, c \in A$.
- A **Riesz space** V is an \mathbb{R} -vector space equipped with a lattice order such that it is an ℓ -group and $0 \leq r$ and $0 \leq v$ imply $rv \geq 0$ for each $r \in \mathbb{R}$ and $v \in V$.

ℓ -groups and Riesz spaces can be axiomatized by equations, and so they form varieties.

Definition

- A map between ℓ -groups is an **ℓ -group homomorphism** if it is a group and a lattice homomorphism.
- An ℓ -group homomorphism between Riesz spaces is a **Riesz space homomorphism** if it is a linear map.

Examples of Riesz spaces



- \mathbb{R}
- \mathbb{R}^X for a set X
- $\mathbb{R} \overrightarrow{\times} \mathbb{R}$ (lexicographic product)
- $C(X, \mathbb{R})$ for a topological space X
- $L^p(\mathbb{R}^n)$

Examples of ℓ -groups

- All the examples above
- \mathbb{Z}
- \mathbb{Z}^X for a set X
- $\mathbb{Z} \overrightarrow{\times} \mathbb{Z}$
- \mathbb{Q}



Definition

- A continuous function $f: \mathbb{R}^k \rightarrow \mathbb{R}$ is **piecewise linear (homogeneous)** if there exist $g_1, \dots, g_n: \mathbb{R}^k \rightarrow \mathbb{R}$ linear homogeneous functions (each in finitely many variables) such that for each $x \in \mathbb{R}^k$ we have $f(x) = g_i(x)$ for some $i = 1, \dots, n$.
- We say that a piecewise linear function f **has integer coefficients**, if it is defined by g_1, \dots, g_n with integer coefficients.



Definition

- A continuous function $f: \mathbb{R}^k \rightarrow \mathbb{R}$ is **piecewise linear (homogeneous)** if there exist $g_1, \dots, g_n: \mathbb{R}^k \rightarrow \mathbb{R}$ linear homogeneous functions (each in finitely many variables) such that for each $x \in \mathbb{R}^k$ we have $f(x) = g_i(x)$ for some $i = 1, \dots, n$.
- We say that a piecewise linear function f **has integer coefficients**, if it is defined by g_1, \dots, g_n with integer coefficients.



We denote by

- $\text{PWL}(\mathbb{R}^\kappa)$ the Riesz space of piecewise linear functions $f: \mathbb{R}^\kappa \rightarrow \mathbb{R}$;
- $\text{PWL}_{\mathbb{Z}}(\mathbb{R}^\kappa)$ the ℓ -group of piecewise linear functions $f: \mathbb{R}^\kappa \rightarrow \mathbb{R}$ with integer coefficients.

Theorem (Baker 1968)

Let κ be a cardinal number.

- *The free Riesz space on κ generators is isomorphic to $\text{PWL}(\mathbb{R}^\kappa)$.*
- *The free ℓ -group on κ generators is isomorphic to $\text{PWL}_{\mathbb{Z}}(\mathbb{R}^\kappa)$.*
- The element $[t]$ of the free algebra correspond to the piecewise linear function that maps $x \in \mathbb{R}^\kappa$ to $t(x) \in \mathbb{R}$.
- The free generators of the free algebra correspond to the projections maps onto each coordinate.

If $X \subseteq \mathbb{R}^{\kappa}$, we denote

- $\text{PWL}(X) = \{f|_X \text{ with } f \in \text{PWL}(\mathbb{R}^{\kappa})\}$,
- $\text{PWL}_{\mathbb{Z}}(X) = \{f|_X \text{ with } f \in \text{PWL}_{\mathbb{Z}}(\mathbb{R}^{\kappa})\}$.

Which Riesz spaces (ℓ -groups) are isomorphic to $\text{PWL}(X)$ ($\text{PWL}_{\mathbb{Z}}(X)$) for some $X \subseteq \mathbb{R}^{\kappa}$?

Congruences in ℓ -groups and Riesz spaces correspond to ℓ -ideals.

Definition

An ℓ -ideal in a Riesz space (ℓ -group) is a subgroup I that is convex, i.e. $|a| \leq |b|$ and $b \in I$ imply $a \in I$.

ℓ -ideals in Riesz spaces are automatically vector subspaces.

Definition

- A proper ℓ -ideal is called **maximal** if it is maximal wrt inclusion.
- A nontrivial Riesz space (ℓ -group) A is **simple** if $\{0\}$ and A are the only ℓ -ideals of A .
- A Riesz space (ℓ -group) is **semisimple** if the intersection of all its maximal ℓ -ideals is $\{0\}$.

Proposition

- An ℓ -group is simple iff it embeds into \mathbb{R} .
- A Riesz space is simple iff it is isomorphic to \mathbb{R} .
- An ℓ -group is semisimple iff it can be subdirectly embedded into a product of sub- ℓ -groups of \mathbb{R} .
- A Riesz space is semisimple iff it can be subdirectly embedded into a power of \mathbb{R} .

Examples

- $\mathbb{R} \overrightarrow{\times} \mathbb{R}$ and $\mathbb{Z} \overrightarrow{\times} \mathbb{Z}$ with the lexicographic order are not semisimple (and hence not simple).
- \mathbb{R} is simple as a ℓ -group and as a Riesz space.
- \mathbb{Z} and \mathbb{Q} are simple ℓ -groups.
- $C(X, \mathbb{R})$ is semisimple for any topological space X .
- $\text{PWL}(X)$ and $\text{PWL}_{\mathbb{Z}}(X)$ are semisimple for any $X \subseteq \mathbb{R}^{\kappa}$.

Theorem (Baker 1968)



- Every semisimple Riesz space is isomorphic to $\text{PWL}(X)$ for some $X \subseteq \mathbb{R}^{\kappa}$.
- Every semisimple ℓ -group is isomorphic to $\text{PWL}_{\mathbb{Z}}(X)$ for some $X \subseteq \mathbb{R}^{\kappa}$.

Theorem (Baker 1968)

- Every semisimple Riesz space is isomorphic to $\text{PWL}(C)$ for some closed cone $C \subseteq \mathbb{R}^k$.
- Every semisimple ℓ -group is isomorphic to $\text{PWL}_{\mathbb{Z}}(C)$ for some closed cone $C \subseteq \mathbb{R}^k$.

Definition

A nonempty subset $C \subseteq \mathbb{R}^k$ is a **closed cone** if it is closed under multiplication by nonnegative scalars and it is closed in \mathbb{R}^k with the euclidean topology.



Let \mathcal{F}_κ be the free Riesz space (ℓ -group) over κ generators.
For any $T \subseteq \mathcal{F}_\kappa$ and $S \subseteq \mathbb{R}^\kappa$, we define the following operators.

$$V(T) = \{x \in \mathbb{R}^\kappa \mid t(x) = 0 \text{ for all } [t] \in T\}$$
$$I(S) = \{[t] \in \mathcal{F}_\kappa \mid t(x) = 0 \text{ for all } x \in S\}.$$

Galois connection

$$T \subseteq I(S) \quad \text{iff} \quad S \subseteq V(T).$$

- $V(T)$ is always a closed cone of \mathbb{R}^κ .
- $I(S)$ is always an ℓ -ideal of \mathcal{F}_κ .

What are the fixpoints of the Galois connection?

$S = VI(S)$ iff S is a closed cone in \mathbb{R}^κ .

S is a fixpoint iff $S = V(T)$ for some $T \subseteq \mathcal{F}_\kappa \cong \text{PWL}(\mathbb{R}^\kappa)$. It can be shown that closed cones are exactly the vanishing sets of families of piecewise linear functions (with integer coefficients) on \mathbb{R}^κ .

$T = IV(T)$ iff T is a ℓ -ideal of \mathcal{F}_κ that is intersection of maximal ℓ -ideals.

T is a fixpoint iff $T = I(S)$ for some $S \subseteq \mathbb{R}^\kappa$ iff $T = \bigcap \{I(x) \mid x \in S\}$. The proper ℓ -ideals of the form $I(x)$ for some $x \in \mathbb{R}^\kappa$ are exactly the maximal ideals of \mathcal{F}_κ (follows from the characterization of simple algebras).

Proposition

The poset of ℓ -ideals of \mathcal{F}_κ that are intersections of maximal ℓ -ideals is dually isomorphic to the poset of closed cones in \mathbb{R}^κ .

We can extend this dual isomorphism to a dual equivalence of categories between the category of semisimple Riesz spaces (ℓ -groups) and the category of closed cones.

On objects:

Let A be a semisimple Riesz space (ℓ -group), then $A \cong \mathcal{F}_\kappa / J$, where J is an intersection of maximal ℓ -ideals of \mathcal{F}_κ . Then map

$$A \mapsto V(J),$$

where $V(J)$ is a closed cone in \mathbb{R}^κ .

Let C be a closed cone in \mathbb{R}^κ . Then map

$$C \mapsto \text{PWL}(C),$$

which is semisimple and isomorphic to $\mathcal{F}_\kappa / I(C)$. (In the case of ℓ -groups map C to $\text{PWL}_{\mathbb{Z}}(C)$.)

On morphisms:

Let $h: A \rightarrow B$ be a Riesz space (ℓ -group) homomorphism with $A \cong \mathcal{F}_\kappa / J_A$ and $B \cong \mathcal{F}_\mu / J_B$. Then map

$$h \mapsto f_h,$$

with $f_h: V(J_B) \rightarrow V(J_A)$ the piecewise linear map whose i^{th} component is given by $h([a_i]) \in \mathcal{F}_\mu / J_B$ where a_i is the i^{th} generator of \mathcal{F}_κ .

Let $f: C \rightarrow D$ be a piecewise linear function (with integer coefficients) between closed cones. Then map

$$f \rightarrow h_f,$$


where $h_f: \text{PWL}(D) \rightarrow \text{PWL}(C)$ is given by $h_f(g) = g \circ f$ (in the case of ℓ -groups we have $h_f: \text{PWL}_{\mathbb{Z}}(D) \rightarrow \text{PWL}_{\mathbb{Z}}(C)$).

These functors yield the Baker-Beynon duality:

Theorem (Beynon 1974)


- The category of *semisimple Riesz spaces* is dually equivalent to the category of *closed cones in \mathbb{R}^k* and piecewise linear maps with real coefficients.
- The category of *semisimple ℓ -groups* is dually equivalent to the category of *closed cones in \mathbb{R}^k* and piecewise linear maps with integer coefficients.

\mathbb{R} (as a Riesz space) is dual to the semiline $\{x \in \mathbb{R} \mid x \geq 0\}$.




\mathbb{R} (as a Riesz space) is dual to the semiline $\{x \in \mathbb{R} \mid x \geq 0\}$.

Indeed, $\mathbb{R} \cong \text{PWL}(\{x \in \mathbb{R} \mid x \geq 0\})$.



$\mathcal{F}_2 / \langle (x - y) \wedge y \wedge 0 \rangle$ is dual to $\{(x, y) \in \mathbb{R}^2 \mid 0 \leq y \leq x\}$.



Generalizing Baker-Beynon duality beyond semisimplicity

In the definition of the operators

$$V(T) = \{x \in \mathbb{R}^\kappa \mid t(x) = 0 \text{ for all } [t] \in T\} \text{ with } T \subseteq \mathcal{F}_\kappa$$

$$I(S) = \{[t] \in \mathcal{F}_\kappa \mid t(x) = 0 \text{ for all } x \in S\} \text{ with } S \subseteq \mathbb{R}^\kappa.$$

we can replace \mathbb{R} with any Riesz space (ℓ -group) A and still get a Galois connection.

In the definition of the operators

$$V(T) = \{x \in A^\kappa \mid t(x) = 0 \text{ for all } [t] \in T\} \text{ with } T \subseteq \mathcal{F}_\kappa$$
$$I(S) = \{[t] \in \mathcal{F}_\kappa \mid t(x) = 0 \text{ for all } x \in S\} \text{ with } S \subseteq A^\kappa.$$

we can replace \mathbb{R} with any Riesz space (ℓ -group) A and still get a Galois connection.

Caramello, Marra, and Spada (2021) observed that this can be done for any variety of algebras by replacing \mathbb{R} with any algebra in that variety. They also show that this approach also works in a more categorical setting.

Our goal is to replace \mathbb{R} with a Riesz space that guarantees more ℓ -ideals of \mathcal{F}_κ to be fixpoints of IV . In this way we extend Baker-Beynon duality beyond semisimple Riesz spaces and ℓ -groups.

It is not possible to obtain a Riesz space (ℓ -group) A such that for any κ the fixpoints of IV are all the ℓ -ideals of \mathcal{F}_κ . This is a consequence of the fact that there are subdirectly irreducible Riesz spaces (ℓ -groups) of arbitrarily large cardinality.

However, if we fix a cardinal α , we will see that we can find A such that for any $\kappa < \alpha$ the fixpoints of IV are all the ideals of \mathcal{F}_κ .

We will see how this yields a duality for all Riesz spaces (ℓ -groups) that are κ -generated (i.e. generated by a set of cardinality at most κ) with $\kappa < \alpha$. In particular, we obtain a duality for all finitely generated Riesz spaces (ℓ -groups) by taking $\alpha = \omega$.

We will replace maximal ℓ -ideals with prime ℓ -ideals.

Definition

An ℓ -ideal I is **prime** if $a \wedge b \in I$ implies $a \in I$ or $b \in I$.

Theorem

- A/I is linearly ordered iff I is prime.
- Every ℓ -ideal is intersection of prime ℓ -ideals.
- Every Riesz space (ℓ -group) is subdirect product of linearly ordered ones.

We fix a cardinal α and we look for a Riesz space (ℓ -group) A into which all the κ -generated with $\kappa < \alpha$ linearly ordered Riesz spaces (ℓ -groups) embed.

Theorem (C., Lapenta, Spada)

Let α be a cardinal. There exists an ultrapower \mathcal{U} of \mathbb{R} in which all κ -generated (with $\kappa < \alpha$) linearly ordered Riesz spaces and ℓ -groups embed.

Proof sketch.

- The theory of nontrivial linearly ordered Riesz spaces is complete. So, each lin. ordered Riesz space $A \neq 0$ is elementarily equivalent to \mathbb{R} .
- Thus, for any cardinal β there is an ultrapower of \mathbb{R} into which all the linearly ordered Riesz spaces of cardinality less than β embed.
- Since a Riesz space that is κ -generated has cardinality at most $\max(\kappa, 2^\omega)$, it is sufficient to take $\beta = \max(\alpha, 2^\omega)$. □

For $\alpha = \omega$ we have can pick \mathcal{U} as follows:

Proposition

Let \mathcal{U} be any ultrapower of \mathbb{R} over a nonprincipal ultrafilter of a countably infinite set. Then every finitely generated linearly ordered Riesz space and ℓ -group embeds into \mathcal{U} .

Fix a cardinal α and \mathcal{U} an ultrapower of \mathbb{R} in which all κ -generated with $\kappa < \alpha$ linearly ordered Riesz spaces and ℓ -groups embed. κ will denote an arbitrary cardinal smaller than α .

We consider the operators:

$$V(T) = \{x \in \mathcal{U}^\kappa \mid t(x) = 0 \text{ for all } [t] \in T\} \text{ with } T \subseteq \mathcal{F}_\kappa$$

$$I(S) = \{[t] \in \mathcal{F}_\kappa \mid t(x) = 0 \text{ for all } x \in S\} \text{ with } S \subseteq \mathcal{U}^\kappa.$$

Galois connection

$$T \subseteq I(S) \quad \text{iff} \quad S \subseteq V(T).$$

- $T = IV(T)$ iff T is an ℓ -ideal of \mathcal{F}_κ .
- We call $S \subseteq \mathcal{U}^\kappa$ such that $S = VI(S)$ a **generalized closed cone** (**\mathbb{Z} -generalized closed cone**).

Proposition

The poset of ℓ -ideals of \mathcal{F}_κ is dually isomorphic to the poset of generalized closed cones (\mathbb{Z} -generalized closed cones) in \mathcal{U}^κ .

Definition

- We say that a map $\mathcal{U}^\kappa \rightarrow \mathcal{U}^\mu$ is **definable** (**\mathbb{Z} -definable**) if its components are defined by terms in the language of Riesz spaces (ℓ -groups).
- If $X \subseteq \mathcal{U}^\kappa$, we denote by $\text{Def}(X)$ and $\text{Def}_{\mathbb{Z}}(X)$ the sets of definable maps and \mathbb{Z} -definable maps $f: X \rightarrow \mathcal{U}$.

The functors $A \cong \mathcal{F}_\kappa / J \mapsto V(J)$ and $C \mapsto \text{Def}(C) \cong \mathcal{F}_\kappa / I(C)$ induce:

Theorem (C., Lapenta, Spada)

- *The category of κ -generated Riesz spaces (with $\kappa < \alpha$) is dually equivalent to the category of **generalized closed cones in \mathcal{U}^κ** (with $\kappa < \alpha$) and definable maps.*

Definition

- We say that a map $\mathcal{U}^\kappa \rightarrow \mathcal{U}^\mu$ is **definable** (\mathbb{Z} -definable) if its components are defined by terms in the language of Riesz spaces (ℓ -groups).
- If $X \subseteq \mathcal{U}^\kappa$, we denote by $\text{Def}(X)$ and $\text{Def}_{\mathbb{Z}}(X)$ the sets of definable maps and \mathbb{Z} -definable maps $f: X \rightarrow \mathcal{U}$.

The functors $A \cong \mathcal{F}_\kappa / J \mapsto V(J)$ and $C \mapsto \text{Def}_{\mathbb{Z}}(C) \cong \mathcal{F}_\kappa / I(C)$ induce:

Theorem (C., Lapenta, Spada)

- *The category of κ -generated Riesz spaces (with $\kappa < \alpha$) is dually equivalent to the category of generalized closed cones in \mathcal{U}^κ (with $\kappa < \alpha$) and definable maps.*
- *The category of κ -generated ℓ -groups (with $\kappa < \alpha$) is dually equivalent to the category of \mathbb{Z} -generalized closed cones in \mathcal{U}^κ (with $\kappa < \alpha$) and \mathbb{Z} -definable maps.*

Consequences and applications of the duality

Proposition

- The generalized closed cones in \mathcal{U}^{κ} (together with \emptyset) form the closed of a topology on \mathcal{U}^{κ} . The closure of a nonempty $X \subseteq \mathcal{U}^{\kappa}$ is $\vee I(X)$.
- \mathbb{R}^{κ} is a subset of \mathcal{U}^{κ} and the closed subsets of \mathbb{R}^{κ} with the subspace topology are exactly the closed cones (and \emptyset).

We obtain the following correspondences:

\mathcal{F}_{κ}	\mathbb{R}^{κ}	\mathcal{U}^{κ}
maximal ℓ -ideals	half-lines from the origin	closures of points of \mathbb{R}^{κ} (except the origin)
intersections of maximal ℓ -ideals	closed cones	closures of nonempty subsets of \mathbb{R}^{κ}
prime ℓ -ideals		irreducible closed subsets = closures of points of \mathcal{U}^{κ} (except the origin)
ℓ -ideals		generalized closed cones

If A is a Riesz space (ℓ -group), then $\text{Spec}(A) = \{\text{prime } \ell\text{-ideals of } A\}$ is called the **spectrum** of A and is naturally equipped with the **Zariski topology** generated by the closed subsets $\{P \in \text{Spec}(A) \mid a \in P\}$, where a ranges in A .

If P is a prime ℓ -ideal of \mathcal{F}_κ , then $V(P)$ is the closure of a point of \mathcal{U}^κ . Choose one such point $x_P \in \mathcal{U}^\kappa$ for each $P \in \text{Spec}(\mathcal{F}_\kappa)$. Let $\mathcal{E}: \text{Spec}(\mathcal{F}_\kappa) \rightarrow \mathcal{U}^\kappa$ be defined by $\mathcal{E}(P) = x_P$.

Theorem (C., Lapenta, Spada)

- \mathcal{E} is a topological embedding.
- \mathcal{E}^{-1} is a complete lattice isomorphism between $\text{Op}(\mathcal{U}^\kappa \setminus \{O\})$ and $\text{Op}(\text{Spec}(\mathcal{F}_\kappa))$.

The spectrum of each Riesz space (ℓ -group) is a **generalized spectral space**, i.e. it is T_0 , sober, the compact open subsets form a basis, and the intersection of two compact opens is compact.

Theorem (C., Lapenta, Spada)

$\mathcal{U}^\kappa \setminus \{O\}$ is a generalized spectral space.

$\mathcal{E}: \text{Spec}(\mathcal{F}_\kappa) \rightarrow \mathcal{U}^\kappa$ can be thought of as a **coordinatization** of $\text{Spec}(\mathcal{F}_\kappa)$ with coordinates in \mathcal{U} .

By the correspondence theorem, if $A \cong \mathcal{F}_\kappa / J$, then we can think of $\text{Spec}(A)$ as a subspace of $\text{Spec}(\mathcal{F}_\kappa)$. So, \mathcal{E} restricts to an embedding of $\text{Spec}(A)$ into \mathcal{U}^κ whose image is $\mathcal{E}[\text{Spec}(\mathcal{F}_\kappa)] \cap V(J)$.

While the spectrum as a topological space is not sufficient to recover the original Riesz space, the coordinatization is enough:

Theorem (C., Lapenta, Spada)

$A \cong \text{Def}(\mathcal{E}[\text{Spec}(A)])$ for any Riesz space A .

An analogous result holds for ℓ -groups.

In part II we will see how $\mathcal{E}[\text{Spec}(\mathcal{F}_\kappa)]$ looks like when κ is finite.

Definition

- Recall that an Riesz space (ℓ -group) is **semisimple** if the intersection of all its maximal ℓ -ideals is $\{0\}$.
- A Riesz space (ℓ -group) A is called **Archimedean** if for every $a, b \in A$, $a \leq 0$ whenever $na \leq b$ for all $n \in \mathbb{N}$.
- Semisimplicity always implies Archimedeanity.
- Archimedeanity implies semisimplicity in the presence of a strong order-unit (e.g. in the finitely generated setting).

Theorem (C., Lapenta, Spada)

Let A be a Riesz space (ℓ -group) and $C \subseteq \mathcal{U}^\kappa$ its dual generalized closed cone (\mathbb{Z} -generalized closed cone).

A is semisimple iff $C = \text{VI}(C \cap \mathbb{R}^\kappa)$, i.e. C is the closure of $C \cap \mathbb{R}^\kappa$ in \mathcal{U}^κ .

Note that $C \cap \mathbb{R}^\kappa$ is the closed cone in \mathbb{R}^κ corresponding to A under Baker-Beynon duality.

For any natural number n let $\pi_n: \mathcal{U}^\omega \rightarrow \mathcal{U}^{n+1}$ be the map that sends $(x_i)_{i \in \omega}$ to (x_0, x_1, \dots, x_n) .

Theorem (C., Lapenta, Spada)

Let A be an ω -generated Riesz space (ℓ -group) and $C \subseteq \mathcal{U}^\omega$ its dual generalized closed cone (\mathbb{Z} -generalized closed cone).

Then A is archimedean iff

$$C = \bigcap_{n=0}^{\infty} \pi_n^{-1}[\text{VI}(\text{VI}(\pi_n[C]) \cap \mathbb{R}^{n+1})],$$

where the subsets $\pi_n^{-1}[\text{VI}(\text{VI}(\pi_n[C]) \cap \mathbb{R}^{n+1})]$ form a decreasing sequence of generalized closed cones in \mathcal{U}^ω .

When $\kappa > \omega$, the decreasing sequence is substituted by a downdirected family of generalized closed cones in \mathcal{U}^κ .

Part II

Using non-standard tools

Recap of Part I

- We derived for any cardinal α a generalization of Baker-Beynon duality to the categories of all Riesz spaces and all ℓ -groups with less than α generators.
- The idea is to replace \mathbb{R} with a suitable ultrapower \mathcal{U} .
- κ -generated Riesz spaces correspond to generalized closed cones in \mathcal{U}^κ . Whereas, κ -generated ℓ -groups correspond to \mathbb{Z} -generalized closed cones in \mathcal{U}^κ .
- Every Riesz space (ℓ -group) can be represented as the algebra of definable (\mathbb{Z} -definable) functions on its dual.
- \mathcal{U}^κ has naturally two topologies (one relative to Riesz spaces and one to ℓ -groups) whose nonempty closed are the generalized closed cones and the \mathbb{Z} -generalized closed cones. These topologies are strictly connected to the spectrum of the free algebra with the Zariski topology.
- Irreducible closed subsets (=closures of points) of $\mathcal{U}^\kappa \setminus \{0\}$ correspond to prime ideals of the free algebra.

For the rest of the talk we will assume $\alpha = \omega$.

Let also assume that \mathcal{U} is an ultrapower of \mathbb{R} defined as $\mathcal{U} = \mathbb{R}^{\mathbb{N}}/\mathcal{F}$ with \mathcal{F} a nonprincipal ultrafilter of $\mathcal{P}(\mathbb{N})$.

We have seen that \mathcal{U} induces dualities for finitely generated Riesz spaces and ℓ -groups.

Theorem

- *The category of all finitely generated Riesz spaces is dually equivalent to the category of generalized closed cones in \mathcal{U}^n (with $n \in \mathbb{N}$).*
- *The category of all finitely generated ℓ -groups is dually equivalent to the category of \mathbb{Z} -generalized closed cones in \mathcal{U}^n (with $n \in \mathbb{N}$).*

It follows from Łoś's theorem that the algebraic structure of \mathbb{R} lifts to \mathcal{U} :

Proposition

- \mathcal{U} is a linearly ordered field.
- \mathcal{U}^n is a \mathcal{U} -vector space.

The elements of \mathcal{U} are equivalence classes $[(r_i)_{i \in \mathbb{N}}]$ of \mathbb{N} -indexed sequences $(r_i)_{i \in \mathbb{N}}$ of real numbers. Where

$$(r_i)_{i \in \mathbb{N}} \sim (s_i)_{i \in \mathbb{N}} \quad \text{iff} \quad \{i \in \mathbb{N} \mid r_i = s_i\} \in \mathcal{F}.$$

We identify each $r \in \mathbb{R}$ with $[(r_i)_{i \in \mathbb{N}}] \in \mathcal{U}$ such that $r_i = r$ for all $i \in \mathbb{N}$.

Proposition

- \mathbb{R} embeds into \mathcal{U} as a sub-lattice-ordered field.
- \mathcal{U}^n is an \mathbb{R} -vector space containing \mathbb{R}^n as a vector subspace.

We will identify \mathbb{R} and \mathbb{R}^n with their isomorphic copies in \mathcal{U} and \mathcal{U}^n .

Some notions from non-standard analysis

As it is common in non-standard analysis, we call the elements of \mathcal{U} **hyperreal numbers**. Among the hyperreal numbers we have:

- **real numbers**

$$[(1, 1, 1, \dots)], \quad \left[\left(\frac{15}{7}, \frac{15}{7}, \frac{15}{7}, \dots \right) \right], \quad [(\pi, \pi, \pi, \dots)], \dots$$

- **infinitesimal numbers** (absolute value smaller than any $0 < r \in \mathbb{R}$)

$$\left[\left(1, \frac{1}{2}, \frac{1}{3}, \dots \right) \right], \quad \left[\left(1, \frac{1}{2^2}, \frac{1}{3^2}, \dots \right) \right], \quad \left[\left(1, \frac{1}{2^2}, \frac{1}{2^3}, \dots \right) \right], \dots$$

- **unlimited numbers** (absolute value greater than any $r \in \mathbb{R}$)

$$[(1, 2, 3, \dots)], \quad [(1, 2^2, 3^2, \dots)], \quad [(1, 2^2, 2^3, \dots)], \dots$$

- **limited numbers** (not limited, i.e. between $-r$ and r for some $r \in \mathbb{R}$)

$$[(1, 1, 1, \dots)], \quad \left[\left(1, \frac{1}{2}, \frac{1}{3}, \dots \right) \right], \quad \left[\left(\frac{1}{2}, \frac{2}{3}, \frac{3}{4}, \dots, 1 - \frac{1}{n}, \dots \right) \right], \dots$$

The operations behave like the limits in analysis:

- limited + limited = limited, unlimited + limited = unlimited, ...
- limited \times limited = limited, unlimited \times infinitesimal = ?, ...

Definition

- If $A \subseteq \mathbb{R}^n$, its **enlargement** ${}^*A \subseteq \mathcal{U}^n$ is defined as follows:

$$\left([(r_i^1)], \dots, [(r_i^n)] \right) \in {}^*A \text{ if and only if } \{i \in \mathbb{N} \mid (r_i^1, \dots, r_i^n) \in A\} \in \mathcal{F}.$$

- If $A \subseteq \mathbb{R}^n$ and $f: A \rightarrow \mathbb{R}$, then the **enlargement** ${}^*f: {}^*A \rightarrow \mathcal{U}$ of f is given by

$${}^*f\left([(r_i^1)], \dots, [(r_i^n)]\right) := [(f(r_i^1, \dots, r_i^n))].$$

Proposition

- $A \subseteq {}^*A$.
- *If A is finite, then $A = {}^*A$.*
- *If A is infinite, then *A must contain some elements of \mathcal{U}^n outside \mathbb{R}^n .*

For example, ${}^*\mathbb{N}$ contains the unlimited element $[(1, 2, 3, \dots)]$.

Let \mathcal{L} be a first-order language and $(\mathbb{R}, (P_\alpha), (f_\alpha))$ an \mathcal{L} -structure, where the P_α 's and f_α 's are the interpretations of the predicate and function symbols of \mathcal{L} in \mathbb{R} . Then $(\mathcal{U}, (*P_\alpha), (*f_\alpha))$ is also an \mathcal{L} -structure.

Theorem (Transfer principle)

*Let φ be a first-order \mathcal{L} -sentence. Then φ is true in $(\mathbb{R}, (P_\alpha), (f_\alpha))$ if and only if φ is true in $(\mathcal{U}, (*P_\alpha), (*f_\alpha))$.*

In other words, a first-order condition holds in \mathbb{R} iff the condition obtained by replacing all the relations and functions with their enlargements holds in \mathcal{U} . (For simplicity of notation, we just write $+$ instead of $*+$ and similarly for the other lattice-ordered field operations.)

This allows to transfer first-order properties of functions and subsets from \mathbb{R}^n to \mathcal{U}^n and back.

Let $S^1 = \{(x, y) \in \mathbb{R}^2 \mid x^2 + y^2 = 1\}$ be the unit circle in \mathbb{R}^2 .
Since



$$\forall x, y ((x, y) \in S^1 \Leftrightarrow x^2 + y^2 = 1)$$

is a first-order condition that holds in \mathbb{R} , then

$$\forall x, y ((x, y) \in {}^*(S^1) \Leftrightarrow x^2 + y^2 = 1)$$


holds in \mathcal{U} by transfer. So, ${}^*(S^1) = \{(x, y) \in \mathcal{U}^2 \mid x^2 + y^2 = 1\}$.

It is easy to get a geometric intuition of the enlargements of subsets of \mathbb{R}^n defined by first-order sentences.




If $0 < \varepsilon \in \mathcal{U}$ is infinitesimal, then $x = \left(\frac{1}{\sqrt{1 + \varepsilon^2}}, \frac{\varepsilon}{\sqrt{1 + \varepsilon^2}} \right) \in {}^*(S^1) \setminus S^1$.


If $0 < \varepsilon \in \mathcal{U}$ is infinitesimal, then $x = \left(\frac{1}{\sqrt{1 + \varepsilon^2}}, \frac{\varepsilon}{\sqrt{1 + \varepsilon^2}} \right) \in {}^*(S^1) \setminus S^1$.





If $0 < \varepsilon \in \mathcal{U}$ is infinitesimal, then $x = \left(\frac{1}{\sqrt{1 + \varepsilon^2}}, \frac{\varepsilon}{\sqrt{1 + \varepsilon^2}} \right) \in {}^*(S^1) \setminus S^1$.



If $0 < \varepsilon \in \mathcal{U}$ is infinitesimal, then $x = \left(\frac{1}{\sqrt{1 + \varepsilon^2}}, \frac{\varepsilon}{\sqrt{1 + \varepsilon^2}} \right) \in {}^*(S^1) \setminus S^1$.



If $f: \mathbb{R}^n \rightarrow \mathbb{R}$ is a function, then the graph of ${}^*f: \mathcal{U}^n \rightarrow \mathcal{U}$ is just the enlargement of the graph of f .



The enlargement of f can be used to compute limits. For example,

$$\lim_{x \rightarrow 0} f(x) = 0 \Leftrightarrow {}^*f(x) \text{ infinitesimal for all } x \text{ infinitesimal.}$$

Definable maps and piecewise linear functions

Let $g: \mathcal{U}^n \rightarrow \mathcal{U}$ be definable, i.e. there is a term t such that $g(x) = t(x)$ for all $x \in \mathcal{U}^n$.


If $f: \mathbb{R}^n \rightarrow \mathbb{R}$ is the piecewise linear function defined by the same term, i.e. $f(x) = t(x)$ for all $x \in \mathbb{R}^n$, then the transfer principle yields

$$\forall x \in \mathbb{R}^n (f(x) = t(x)) \quad \text{iff} \quad \forall x \in \mathcal{U}^n (*f(x) = t(x)).$$


Thus, $g = *f$, and so g is the enlargement of a piecewise linear function.

Proposition

Let $C \subseteq \mathcal{U}^n$ be a generalized closed cone. Then $\text{Def}(C) = \{(*f)|_C \mid f: \mathbb{R}^n \rightarrow \mathbb{R} \text{ piecewise linear}\}$.



$$f: \mathbb{R}^2 \rightarrow \mathbb{R}$$




$$*f: \mathcal{U}^2 \rightarrow \mathcal{U}$$

Definable functions naturally generalize piecewise linear functions.

Let $\mathbb{R} \overrightarrow{\times} \mathbb{R}$. Then its dual generalized closed cone is

$$C = \{(x, y) \in \mathcal{U}^2 \mid x > 0, y \geq 0, \text{ and } y/x \text{ is infinitesimal}\} \cup \{(0, 0)\}.$$



So,

$$\begin{aligned} \mathbb{R} \overrightarrow{\times} \mathbb{R} &\cong \text{Def}(C) = \{(*f)|_C \mid f: \mathbb{R}^2 \rightarrow \mathbb{R} \text{ piecewise linear}\} \\ &= \{(*f)|_C \mid f: \mathbb{R}^2 \rightarrow \mathbb{R} \text{ linear}\}. \end{aligned}$$

Indexes and irreducible closed subsets

Recall from part I that f.g. linearly ordered Riesz spaces correspond to the irreducible closed subsets of \mathcal{U}^n , i.e. the closures of the points of \mathcal{U}^n .

We want to understand how these subsets of \mathcal{U}^n look like (for simplicity we only consider the case of Riesz spaces).

Theorem (Orthogonal decomposition)

If $x \in \mathcal{U}^n$, then $x = \alpha_1 v_1 + \dots + \alpha_k v_k$ where $\alpha_1, \dots, \alpha_k \in \mathcal{U}$ are positive, α_{i+1}/α_i is infinitesimal for each $i < k$, and $v_1, \dots, v_k \in \mathbb{R}^n$ are orthonormal vectors. Furthermore, this decomposition is unique.

Definition

- We call a finite sequence (v_1, \dots, v_k) of orthonormal vectors in \mathbb{R}^n an **index**.
- We denote by $\iota(x)$ the index (v_1, \dots, v_k) made of the vectors appearing in the orthogonal decomposition of $x \in \mathcal{U}^n$.
- Let \mathbf{v}, \mathbf{w} be two indexes. We write $\mathbf{v} \leq \mathbf{w}$ when \mathbf{v} is a truncation of \mathbf{w} , i.e. $\mathbf{v} = (v_1, \dots, v_h)$ and $\mathbf{w} = (v_1, \dots, v_k)$ for $h \leq k$.

Definition

If \mathbf{v} is an index, let $\text{Cone}(\mathbf{v}) := \{y \in \mathcal{U}^n \mid \iota(y) \leq \mathbf{v}\}$

Theorem (C., Lapenta, Spada)

The closure of x in \mathcal{U}^n is $\text{Cone}(\iota(x))$.

The proof uses the fact that if $f: \mathbb{R}^n \rightarrow \mathbb{R}$ is a linear function and $x \in \mathcal{U}^n$ with $\iota(x) = (v_1, \dots, v_k)$, then the sign of ${}^*f(x)$ is determined by the real numbers $f(v_1), \dots, f(v_k)$.

Corollary


If $x \in \mathcal{U}^n$, then

$$\begin{aligned} \text{Def}(\text{Cone}(\iota(x))) &\cong \{{}^*f(x) \in \mathcal{U} \mid f: \mathbb{R}^n \rightarrow \mathbb{R} \text{ piecewise linear}\} \\ &= \{{}^*f(x) \in \mathcal{U} \mid f: \mathbb{R}^n \rightarrow \mathbb{R} \text{ linear}\}. \end{aligned}$$

Let $x = (1, 0) \in \mathcal{U}^2$. Then $\iota(x) = (v_1)$ with $v_1 = (1, 0)$. We have

$$y \in \text{Cone}(\iota(x)) \quad \text{iff} \quad y = \alpha_1(1, 0) \text{ with } 0 \leq \alpha_1 \in \mathcal{U}.$$

Thus, the closure of x in \mathcal{U}^2 is $\{(\alpha_1, 0) \mid 0 \leq \alpha_1 \in \mathcal{U}\}$, which is the enlargement of the positive x -semiaxis.



The dual Riesz space is \mathbb{R} . Indeed,

$$\text{Def}(\text{Cone}(\iota(x))) \cong \{^*f(1, 0) \mid f: \mathbb{R}^2 \rightarrow \mathbb{R} \text{ linear}\} \cong \mathbb{R}.$$

Let $\varepsilon \in \mathcal{U}$ be a positive infinitesimal and $x = (1, \varepsilon)$. Then


$$x = 1(1, 0) + \varepsilon(0, 1)$$

is the orthogonal decomposition of x . Thus, $\iota(x) = (v_1, v_2)$ with $v_1 = (1, 0)$ and $v_2 = (0, 1)$. We have

$$y \in \text{Cone}(\iota(x)) \quad \text{iff} \quad y = O, \text{ or} \\ y = \alpha_1(1, 0) \text{ (orthogonal decomposition), or} \\ y = \alpha_1(1, 0) + \alpha_2(0, 1) \text{ (orthogonal decomposition)}$$

Then $\text{Cone}(\iota(x))$, i.e. the closure of x in \mathcal{U}^2 is

$$\{(\alpha_1, \alpha_2) \in \mathcal{U}^2 \mid \alpha_1 > 0, \alpha_2 \geq 0 \text{ and } \alpha_2/\alpha_1 \text{ is infinitesimal}\} \cup \{O\}.$$



The dual Riesz space is $\mathbb{R} \overrightarrow{\times} \mathbb{R}$. Indeed,

$$\text{Def}(\text{Cone}(\iota(x))) \cong \{*f(1, \varepsilon) \mid f: \mathbb{R}^n \rightarrow \mathbb{R} \text{ linear}\} \\ = \{a + b\varepsilon \in \mathcal{U} \mid a, b \in \mathbb{R}\} \cong \mathbb{R} \overrightarrow{\times} \mathbb{R}.$$

Theorem (C., Lapenta, Spada)

The mapping $\text{Cone}: \mathbf{v} \mapsto \text{Cone}(\mathbf{v})$ induces an order-isomorphism between the set of indexes ordered by truncation and the set of irreducible closed subsets of \mathcal{U}^n ordered by inclusion.

Corollary

$\text{I} \circ \text{Cone}: \mathbf{v} \mapsto \text{I}(\text{Cone}(\mathbf{v}))$ induces an order-isomorphism between the set of nonempty indexes ordered by truncation and $\text{Spec}(\mathcal{F}_n)$ ordered by reverse inclusion.

That nonempty indexes correspond to prime ideals of \mathcal{F}_n was proved by [Panti \(1999\)](#) using different techniques.

Embedding $\text{Spec}(\mathcal{F}_n)$ into \mathcal{U}^n

Recall from part I: if we choose for each irreducible closed subset $C \subseteq \mathcal{U}^n \setminus \{O\}$ a point $x \in \mathcal{U}^n$ such that C is the closure of x , then we can define an embedding $\mathcal{E}: \text{Spec}(\mathcal{F}_n) \rightarrow \mathcal{U}^n$.


Indexes allow us to choose x for every C in a canonical way. Fix a positive infinitesimal $\varepsilon \in \mathcal{U}$. If $C = \text{Cone}(\mathbf{v})$ is an irreducible closed with $\mathbf{v} = (v_1, \dots, v_k)$, then we pick $x \in \text{Cone}(\mathbf{v})$ defined as

$$x = v_1 + \varepsilon v_2 + \dots + \varepsilon^{k-1} v_k.$$

Since $\mathbf{v} = \iota(x)$, we have that $\text{Cone}(\mathbf{v})$ is the closure of x .


Therefore, we obtain an embedding $\mathcal{E}: \text{Spec}(\mathcal{F}_n) \rightarrow \mathcal{U}^n$ that maps a prime ideal $P = \mathfrak{l}(\text{Cone}(\mathbf{v}))$ to the point $v_1 + \varepsilon v_2 + \dots + \varepsilon^{k-1} v_k$.

We have $\mathcal{E}[\text{Spec}(\mathcal{F}_1)] = \{-1, 1\} \subseteq \mathcal{U}$.




Note that $\text{Spec}(\mathcal{F}_1) = \text{MaxSpec}(\mathcal{F}_1)$.


We have $\mathcal{E}[\text{MaxSpec}(\mathcal{F}_2)] = S^1 \subseteq \mathbb{R}^2 \subseteq \mathcal{U}^2$.



We have $\mathcal{E}[\text{Spec}(\mathcal{F}_2)] \subseteq \mathcal{U}^2$ consists of points infinitesimally close to S^1 .




We have $\mathcal{E}[\text{Spec}(\mathcal{F}_2)] \subseteq \mathcal{U}^2$ consists of points infinitesimally close to S^1 .




$\text{Spec}(\mathcal{F}_2)$

We have $\mathcal{E}[\text{Spec}(\mathcal{F}_2)] \subseteq \mathcal{U}^2$ consists of points infinitesimally close to S^1 .




$\text{Spec}(\mathcal{F}_2)$


We have $\mathcal{E}[\text{Spec}(\mathcal{F}_2)] \subseteq \mathcal{U}^2$ consists of points infinitesimally close to S^1 .




We have $\mathcal{E}[\text{MaxSpec}(\mathcal{F}_3)] = S^2 \subseteq \mathbb{R}^3 \subseteq \mathcal{U}^3$.



We have $\mathcal{E}[\text{Spec}(\mathcal{F}_3)] \subseteq \mathcal{U}^3$ consists of points infinitesimally close to S^2 .




We have $\mathcal{E}[\text{Spec}(\mathcal{F}_3)] \subseteq \mathcal{U}^3$ consists of points infinitesimally close to S^2 .



$\text{Spec}(\mathcal{F}_3)$

We have $\mathcal{E}[\text{Spec}(\mathcal{F}_3)] \subseteq \mathcal{U}^3$ consists of points infinitesimally close to S^2 .



Characterization of prime ideals in \mathcal{F}_n

We have seen that $I \circ \text{Cone}$ induces an order-isomorphism between indexes and prime ideals of \mathcal{F}_n . Recall that $\mathcal{F}_n \cong \text{PWL}(\mathbb{R}^n)$

$I(\text{Cone}(\mathbf{v}))$ correspond to the prime ideal of $\text{PWL}(\mathbb{R}^n)$ given by

$$\{f \in \text{PWL}(\mathbb{R}^n) \mid *f \text{ vanishes on } \text{Cone}(\mathbf{v})\}.$$

Is there a way to avoid mentioning the enlargement?

Definition


Let $\mathbf{v} = (v_1, \dots, v_k)$ be an index. We say that a closed cone $C \subseteq \mathbb{R}^n$ is a \mathbf{v} -cone if there exist real numbers $0 < r_1, \dots, r_k \in \mathbb{R}$ such that C is the positive span of

$$\{r_1 v_1, \quad r_1 v_1 + r_2 v_2, \quad \dots \quad r_1 v_1 + \dots + r_k v_k\}.$$


Theorem (C., Lapenta, Spada)

$\text{Cone}(\mathbf{v})$ is the intersection of the enlargements of all \mathbf{v} -cones.


Let $\mathbf{v} = (v_1, v_2)$ with $v_1 = (1, 0)$ and $v_2 = (0, 1)$.




Let $\mathbf{v} = (v_1, v_2)$ with $v_1 = (1, 0)$ and $v_2 = (0, 1)$.




Let $\mathbf{v} = (v_1, v_2)$ with $v_1 = (1, 0)$ and $v_2 = (0, 1)$.



Let $\mathbf{v} = (v_1, v_2)$ with $v_1 = (1, 0)$ and $v_2 = (0, 1)$.




Let $\mathbf{v} = (v_1, v_2)$ with $v_1 = (1, 0)$ and $v_2 = (0, 1)$.




$\bigcap \{C \mid C \text{ is a } \mathbf{v}\text{-cone}\}$ is the positive x -semiaxis.


Let $\mathbf{v} = (v_1, v_2)$ with $v_1 = (1, 0)$ and $v_2 = (0, 1)$.




Let $\mathbf{v} = (v_1, v_2)$ with $v_1 = (1, 0)$ and $v_2 = (0, 1)$.




Let $\mathbf{v} = (v_1, v_2)$ with $v_1 = (1, 0)$ and $v_2 = (0, 1)$.



Let $\mathbf{v} = (v_1, v_2)$ with $v_1 = (1, 0)$ and $v_2 = (0, 1)$.



Let $\mathbf{v} = (v_1, v_2)$ with $v_1 = (1, 0)$ and $v_2 = (0, 1)$.



$$\bigcap \{ *C \mid C \text{ is a } \mathbf{v}\text{-cone} \} = \text{Cone}(\mathbf{v}).$$


Theorem (C., Lapenta, Spada)

*$*f$ vanishes on $\text{Cone}(\mathbf{v})$ iff f vanishes on a \mathbf{v} -cone.*

Proof sketch.

- By transfer, if f vanishes on a \mathbf{v} -cone C , then $*f$ vanishes on $*C$. So, $*f$ vanishes on $\text{Cone}(\mathbf{v})$ because $\text{Cone}(\mathbf{v}) \subseteq *C$.






Theorem (C., Lapenta, Spada)

f vanishes on Cone(v**) iff f vanishes on a **v**-cone.*

Proof sketch.

- By transfer, if f vanishes on a \mathbf{v} -cone C , then $*f$ vanishes on $*C$. So, $*f$ vanishes on $\text{Cone}(\mathbf{v})$ because $\text{Cone}(\mathbf{v}) \subseteq *C$.
- If $*f$ vanishes on $\text{Cone}(\mathbf{v})$, then there are $0 < \alpha_1, \dots, \alpha_k \in \mathcal{U}$ such that the positive span S of $\{\alpha_1 v_1, \dots, \alpha_1 v_1 + \dots + \alpha_k v_k\}$, is contained in $\text{Cone}(\mathbf{v})$.





Theorem (C., Lapenta, Spada)

**f vanishes on Cone(v) iff f vanishes on a v-cone.*

Proof sketch.

- By transfer, if f vanishes on a \mathbf{v} -cone C , then $*f$ vanishes on $*C$. So, $*f$ vanishes on $\text{Cone}(\mathbf{v})$ because $\text{Cone}(\mathbf{v}) \subseteq *C$.
- If $*f$ vanishes on $\text{Cone}(\mathbf{v})$, then there are $0 < \alpha_1, \dots, \alpha_k \in \mathcal{U}$ such that the positive span S of $\{\alpha_1 v_1, \dots, \alpha_1 v_1 + \dots + \alpha_k v_k\}$, is contained in $\text{Cone}(\mathbf{v})$.
- Thus, $*f$ vanishes on S . By the transfer principle, there are $0 < r_1, \dots, r_k \in \mathbb{R}$ such that f vanishes on the positive span of $\{r_1 v_1, \dots, r_1 v_1 + \dots + r_k v_k\}$, which is a \mathbf{v} -cone.

□

We obtain the characterization of prime ideals of \mathcal{F}_n due to [Panti \(1999\)](#).

Corollary

$I(\text{Cone}(\mathbf{v})) = \{f \in \text{PWL}(\mathbb{R}^n) \mid f \text{ vanishes on a } \mathbf{v}\text{-cone}\}.$

- Extend Marra-Spada duality for semisimple MV-algebras and Riesz MV-algebras beyond semisimplicity:
 - $\mathcal{U} \rightarrow {}^*[0, 1]$.
 - Indexes: $(v_1, \dots, v_k) \rightarrow (x_0, v_1, \dots, v_k)$.
 - Irreducible closed subsets:
“infinitesimally wide cones” \rightarrow “infinitesimal simplexes”.
 - Describe dually the equivalence between MV-algebras and ℓ -groups with strong order-unit.
 - Describe dually the equivalence between perfect MV-algebras and ℓ -groups.
- Generalize the non-standard techniques and the indexes to the infinitely generated case.
- Use the duality to characterize the free Riesz spaces over ℓ -groups.

THANK YOU!

Theorem

Let $\alpha \leq 2^\omega$ be a cardinal. There exists an ultrapower \mathcal{U} of \mathbb{R} such that every linearly ordered Riesz space an ℓ -groups of cardinality less than α embeds \mathcal{U} .

Proof.

- All nontrivial linearly ordered Riesz spaces are elementarily equivalent: their theory has quantifier elimination, and hence it is model complete. Since \mathbb{R} embeds into every non-trivial Riesz space, the theory of linearly ordered Riesz Spaces is complete because it is model complete and has an algebraically prime model.
- By a model-theoretic fact any α -regular ultrapower \mathcal{U} of \mathbb{R} is such that all linearly ordered groups of cardinality less or equal to α embed into \mathcal{U} .
- Since $\alpha \leq 2^\omega$ (the cardinality of the language of Riesz spaces), another model theoretic fact tells us that every ℓ -group of cardinality less than α embeds into \mathcal{U} .



Definition (Panti (1999))

- We call a subspace of \mathbb{R}^n **rational** if it admits a basis made of vectors from \mathbb{Q}^n .
- If $S \subseteq \mathbb{R}^n$, then its **rational envelope** $\langle S \rangle$ denotes the smallest rational subspace of \mathbb{R}^n containing S
- We say that an index $\mathbf{v} = (v_1, \dots, v_k)$ is \mathbb{Z} -reduced if $v_i \in \langle v_j \rangle^\perp$ for any $i \neq j$.

Given an index \mathbf{v} there is a canonical way to associate a \mathbb{Z} -reduced index $\text{red}(\mathbf{v})$.

Theorem (C., Lapenta, Spada)

- *The closure of x in \mathcal{U}^n with the topology of the \mathbb{Z} -generalized closed cones is $\bigcup \{ \text{Cone}(\mathbf{v}) \mid \text{red}(\mathbf{v}) \leq \text{red}(\iota(x)) \}$.*
- *There is an order isomorphism between \mathbb{Z} -reduced indexes and irreducible closed subsets of \mathcal{U}^n with the topology of the \mathbb{Z} -generalized closed cones.*

Definition

Let A be a Riesz space (ℓ -group) and $0 < a \in A$.

- a is a **strong order-unit** if for each $b \in A$ there exists $n \in \mathbb{N}$ such that $b \leq na$.
- a is a **weak order-unit** of A if $a \wedge |b| = 0$ implies $b = 0$ for each $b \in A$.

Theorem

Let A be a nontrivial Riesz space (ℓ -group) and $C \subseteq \mathcal{U}^k$ its dual generalized closed cone (\mathbb{Z} -generalized closed cone).

- A has a strong order-unit iff $C \setminus \{0\}$ is compact.
- A has a weak order-unit iff $C \setminus \{0\}$ contains a dense compact open subset.

Let A, B be two Riesz spaces dual to the generalized closed cones $C \subseteq U^\kappa$ and $D \subseteq U^\mu$.

Theorem

- The coproduct $A \oplus B$ is dual to $C \times D$.
- The product $A \times B$ is dual to $(C \times \{0\}) \cup (\{0\} \times D)$.
- The lexicographic product $\mathbb{R} \overrightarrow{\times} B$ is dual to

$\{(x, y) \in \mathcal{U} \times D \mid 0 < x, y/x \text{ has all infinitesimal coordinates}\} \cup \{0\}$.